Civic Science: Building Tools and Platforms for Rigorous Public Research

48 ppm

Sara Wylie PhD

Assistant Professor of Sociology/Anthropology

And Health Sciences

Northeastern University

Publiclab.org co-founder/organizer

JPB Environmental Health Fellow







All results support the finding that greatest amount of H2S is found in drainage canal, this is scarey because it can mix with irrigation and surface water that waters crops and livestock.

How Civic Science Can Address:

Emerging Health Threats


Health Disparities/CBPR

Data Gaps

Scientific Literacy

Public Lab's Background







Helium balloon with digital camera attached



Grassroots mapping kit


Public Lab started through developing a cheap method of making satellite-like maps using helium balloons and digital cameras. The method was used during the BP gulf oil spill by communities to create a public archive maps documenting damage from the spill.


Publiclab.org

Health Disparities/CBPR


Emerging Health Threats

Mapknitter.org





ERMA GULF RESPONSE









Can we do better?

Jen Hudon



Barriers to Community-based Environmental Science and Health Research

- Expensive
- Limited to hands of experts
- Lag in knowledge between experts and people on the ground
- People on ground don't own the data
- They also don't understand how scientific knowledge is made
- Industries have far greater capacity to generate data



publiclab.org

Forum for Public R+D

Home »

Balloon & Kite Mapping

View What links here Edit Revisions Tags Track



How Can I Do This?

Our whole toolkit is linked out below, but really fast: 1)Buy our balloon kit, assemble your own, or buy/make a kite.

- 2) find a good camera.
- determine how you will trigger the camera, (we suggest a rubber band) or pick out a timelapse app for your smartphone.
- 4)build a simple housing from a plastic bottle.
- 5)find a site to map that is five miles from an airport and not Washington D.C.
- 6) follow the pre-flight checklist(pdf) and quickstart quide(pdf) to safely fill up your balloon and fly!
- 7) sort your images on your desktop or with Mapmill.
- 8) make them into a map with Mapknitter.
- Print a poster of your map from Mapknitter, see your map join the public record in our archive, and if you'd like,



Balloon mapping kits are now available:

Purchase a kit »

Contributors

PLOTS members who have <u>contributed</u> <u>research notes</u> or <u>added to wiki pages</u> on this topic:

- Jeffrey Warren (283)
- Stewart Long (87)
- MicheleTobias

 (7)
- Adam Griffith (10)
- Shannon (23)
- Liz Barry (161)
- Mathew (104)
- Cesar Harada
 (2)
- Jaekyung lee (3)
- Jeremy Crampton (6)
- maning (2)



Maps open data from balloon and kite photography

Do-It-Yourself "satellite" imagery

These maps were largely made by taking photos from balloons and kites, a technique adopted and refined by Public Lab contributors. Make one yourself and it can be featured here.

☐ A grassroots data archive

This archive represents the collective work of our community to provide an alternative source for aerial imagery, and to highlight issues of environmental and social concern with Do-It-Yourself tools. The archive provides:

- A permanent, backed up archive
 A place to advocate around your
- A space to discuss and understand the maps

by TNGIC | 4 months ago | 〇 0 | ☆ 0

State Park. Burns,

Tennessee



Oakland Cemetery.



Rotsoord. Utrecht, Netherlands. by keimpe, Keimpe de He

towards r...

3 months ago | ♀ 0 | ☆ 0

I just set up a new media lab in Utrecht, the Netherlands called Rotslab. We are peared



Crossing, Pennsylvania.

by seankmcginnis | 4 mon ago | ♀ 0 | ☆ 0



Midtown Atlanta. Atlanta, Georgia.

by The Final Five | 3 months ago | 0 0 | 0 0








Georgia Tech TV


create collaborative workflows

Home »

Research notes for balloon-mapping

Quick process notes and field reports





Analytics on aerial Coded ground control points

by: Jasja

1 week 1 day ago

Ground control point targets are useful to stitch photos on uniform or repeatitive landscapes. They... Read more »



3-D Images- Big Branch Balloon Mapping by: astoicof


1 week 2 days ago

After the balloon mapping trip at Big Branch National Wildlife Refuge on May 14th 2012, I created... Read more »

Tags: big-branch, gulf-coast, louisiana




FABAoutfit for vertical KAP and BAP by: Juan



NRG images of cover crop trials by: cfastie

restoration over time; air vs ground

protect openness with viral licensing





Creative Commons
Share Alike with Attribution For Media




CERN Open Hardware 1.0

Share Back with Attribution


GNU Public License 3.0
Copyleft Viral Software Licensing

How can these tools enable scientific rigor?

- 1. Meta-data gathered constitutively: GPS, time/date stamped.
- 2. Tools can be standardized: community use standardized kits and can demonstrate tool calibration.
- 3. Raw data can be shared publicly and modification to the data can be tracked.
- 4. As communities can build their own tools, experiments can be repeated from place to place.
- 5. Can be designed to integrate with and potentially improve the quality of "standard" data formats.



How can this improve science?









Discovery of unmapped spring



News Release



Region 2 - New Jersey, New York, Puerto Rico and the U.S. Virgin Islands

EPA Proposes Plan for Cleaning Up Gowanus Canal

Multi-million Dollar Cleanup to Revitalize Polluted Brooklyn Waterway

Contact: John Martin, (212) 637-3662, martin.johnj@epa.gov

For the first and second segments of the canal, the EPA is proposing to dredge approximately 307,000 cubic yards of highly contaminated sediment. In some areas where the sediment is contaminated with liquid coal tar, the EPA is proposing to stabilize the sediment by mixing it with concrete or similar materials. The stabilized areas would then be covered with multiple layers of clean material, including an "active" layer made of a specific type of clay that will remove PAH contamination that could well up from below, an "isolation" layer of sand and gravel that will ensure that the contaminants are not exposed, and an "armor" layer of heavier gravel and stone to prevent erosion of the underlying layers from boat traffic and currents. Finally, clean sand would be placed on top of the "armor" layer to restore the canal bottom as a habitat. The plan also calls for removing contaminated material placed in the 1st

Street Turning Basin decades ago.




Grassicots Community Data Collection helped focus attention on this site


Civic Science Approach to Environmental Health Monitoring


- Civic vs. Citizen Science
- Tools developed and owned by communities
- Tools can be adapted and spread in an open source fashion
- Research can be spread and scaled between communities
- Supports public exploration and investigation of environments
- Increases the capacity of regulatory agencies

<u>Institutions for civic technoscience: How critical making is transforming environmental research</u>

How can this help Agencies?







over 130,000 assessments by over 6000 online individuals in one week.


Emerging Health Threats

Data Gaps

Health Disparities/CBPR

Growth of Unconventional Oil and Gas

"Between 15 to 25 percent of natural gas in the U.S. may contain hydrogen sulfide, while worldwide, the figure could be as high as 30 percent." (Skrtic, 2006)



Background: What is Hydrogen Sulfide (H₂S)?

- Hazardous, neurotoxic gas
- Smells like rotten eggs
- Natural sources: volcanoes, stagnant bodies of water
- Industrial sources: sewage plants, CAFOs, pulp and paper mills, oil and gas operations
- Eco-toxic to aquatic life at very low concentrations: 0.0149 ppm 0.0448 ppm (EPA 2012)

• Human Health Impacts (Acute): **Health effects**: eye irritation, rashes, headaches to

serious neurological harm, knock down and/or death

(Kilburn, Thrasher and Gray 2010)

NIOSH Recommended Exposure Limit (REL): 10 ppm, 10-minute ceiling

Concentration considered immediately dangerous to life and health (IDLH): 100 ppm

ACGIH[®] recommends a threshold limit value (TLV[®]) of 1 ppm as an 8-hour time weighted average (TWA) and a short-term exposure limit (STEL) of 5 ppm.



(Skrtic 2006)

At least 15.3 million Americans lived within a mile of a well that has been drilled since 2000. That is more people than live in Michigan or New York City. *Wall Street Journal* 2013 Oct 25


Oil and Gas Wells Near Residences, Pavillion Area, Fremont County, WY

Figure: Residents have health issues they fear are from water contamination caused by oil and gas development. Air sample results in the Pavillion area exceed health based standards, causing residents to worry that air emissions are also impacting their health. Our data shows high levels of volatile organic compounds coming from production equipment and produced water tanks near their homes and farms.

Development Site
 Residence
 O.425
 O.85
 I.7 Miles

Author: Caitlin Kennedy, Source: Garmin eTrex Venture, Source: National Atlas, Source: Fremont County Assessor's Office, Source: EPA Pavillion Area Working Group, Data Frame: NA Equidistant and Equal Area Conic, Geographic Projection: NAD 1983, Date Created: Fall 2012 and Winter 2013

Holes in Data: Existing Stationary Monitors are Insufficient



Holes in Data: Brief History of H₂S Regulation

TRI Reporting is not required for downstream oil and gas operations

- 1990 Clean Air Act Passed, H₂S on list of extremely hazardous substances (1 of 2 to be struck)
- 1994 EPA placed an administrative stay on H₂S reporting (lasted 17 years)
- 2012 Reporting for H₂S is required
 But oil and gas industry is exempt from TRI reporting at downstream facilities (wellheads and impoundments).

GAO (2012:35): "If oil and gas exploration and production were added to the industries

required to report to the TRI, such facilities meeting relevant thresholds

would have to report releases of hydrogen sulfide."


No Protection for Residents





Deb Thomas at the Crosby drill pad near her Wyoming home.

(Warning Signs 2014)



Research on chronic exposure to H₂S

- Respiratory distress
- Headaches
- Depression
- Central Nervous System

Disorders

- Fatigue
- Dizziness
- Anxiety
- Sleep problem
- Memory Loss
- Miscarriage
 - Women employed in rayon textile and paper products jobs in Finland were found to have an increased rate of spontaneous abortions when the mean annual level of hydrogen sulfide exceeded 3 ppb (Hemminki and Niemi 1982).
 - An increase in spontaneous abortions was also found in women working in petrochemical plants in China as compared to women working in non-chemical plants (Xu et al. 1998).

			. ;
Chronic Exposure	Parti-Pellinen et al.	Medicine / Public	Examined health effects chronic, low-level exposure to sulfur compounds,
	(1996)	Health	including H2S, near a paper and pulp mill; administered cross-sectional
			questionnaire to 336 subjects and to a reference community; increased
			frequency of eye and nasal symptoms, coughs, and headaches or
	Lagator et al. (2001)	Medicine /	migraines, and acute respiratory infections. Investigate effects of chronic exposure to low levels of hydrogen sulfide;
	Legator et al. (2001)	Toxicology / Public	multi-symptom health survey submitted to two exposed communities –
		Health	Odessa, Texas and Puna, Hawaii, and to control communities; found
		Ticular	central nervous system impacts: fatigue, restlessness, depression, short
			term memory loss, balance and sleep problems, anxiety, lethargy,
			headaches, dizziness, tremors; respiratory system impacts: wheezing,
			shortness of breath, coughing; and various ear, nose, and throat symptoms.
	Tarver and Dasgupta	Chemistry	To determine hydrogen sulfide concentrations near oil fields in Western
	(1997)		Texas
	Xu et al. (1998)	Medicine /	To determine effects of exposure to hydrogen sulfide on the reproductive
		Epidemiology	system; conducted a retrospective epidemiological study to assess the
			association between spontaneous abortion and exposure to petrochemicals
	77'11 (1000)	P 11 11	in Beijing, China; found an association.
	Kilburn (1999)	Epidemiology	To determine long-term effects of exposure to hydrogen sulfide; examined
			and submitted a questionnaire to four groups of people that were exposed to hydrogen sulfide (from boreholes in the ground, downwind of a
			refinery, due to an oil refinery explosion, and a group of people exposed to
			odors); found abnormal balance, delayed verbal recall, impaired color
			discrimination and grip strength.
	Schiffman et al. (1995)	Psychiatry	To determine the effect of persistent environmental odors on the mood of
			people living near the source of odors; submitted a questionnaire to 44
			subjects and 44 controls; found more tension, depression, fatigue, and
	TZ'11 (2002)	D 11 11	confusion, and less vigor among the exposed subjects.
	Kilburn (2003)	Epidemiology	To measure long term effects of hydrogen sulfide exposure – various
			lengths of exposure and various concentrations; submitted a questionnaire, and performed neuropsychological and neurophysiological tests on 19
			exposed subjects and 202 unexposed subjects; found elevated tensions,
			depression, anger, fatigue, and confusion, and more prevalent respiratory
			symptoms among exposed subjects.

(Skrtic 2006)

Present H₂S Monitoring tools are inaccessible for communities:



- High-cost
- Designed for expert use
- Designed for Industry to meet OSHA standards
- Designed for emergency situations
- Acute, high does exposures
- To monitor individual exposures

How does the community exposure experience differ?

H₂S From Community Perspective:

Exposure Experience:

- Low dose
- Chronic exposures
- Community rather than individual exposures
- Environmental and non human health rather than just human health


Social Exposure Experience:

- Agency inattention
- No training about risks
- Absence of data
- Lack of regulations
- Low social and economic capital relative to industry
- Ambivalence: a feeling of "being beaten down"

Photopaper Method:

Could this Empower Communities and call Attention to H₂S Data Gap?

C.J. Horwell et al. / Journal of Volcanology and Geothermal Research 139 (2005) 259-269




- Low cost
- Designed and developed with exposed communities
- Easy to use
- Map large areas effectively
- Easy to include controls
- Results are visually compelling
- Results are readily interpreted
- Sensitive to low levels:
 - .03 ppm-2.5 ppm






H2S Canisters, McMullan Property, Deaver Area, WY




Author: Cait Kennedy Data Frame: NA Equidistant Conic Geographic Projection: NAD 1983 Source: Garmin eTex Device Date: Summer 2013

How Do You Make It?









48ppm



acute exposure threshold .0653 ppm

intermediate exposure threshold .0187 ppm



Sampled levels of H_2S 735 X's greater than the intermediate exposure threshold of the Federal ATSDR Standards

(Macey et. al. 2014)

(Image credit: Megan McLaughlin)

48ppm

Health Disparities/CBPR

Emerging Health Threats

Scientific Literacy


Data Gaps



All results support the finding that greatest amount of H2S is found in drainage canal, this is scarey because it can mix with irrigation and surface water that waters crops and livestock.

(Image credit: Megan McLaughlin)

Publication and Public Lab Results:



Public Lab Tools Currently In development




Balloon & Kite Mapping



Near-Infrared Camera




Thermal photography



Spectrometer Status: In development



Indoor Air Quality Mapping



Kite-Balloon Hybrid



Balloon Telemetry Kit



Stereo Camera Status: In development



Hydrogen Sulfide Sensing



Home Testing for Water Quality **Endocrine** Disruptors



Sensor

Status: Early adopter only



Air Column Monitor

Status: In development


How Can Public Health Laboratories Support Citizen Science?


 Advise on the development and testing of citizen science tools.

 Support community's scientific research and planning.


Act as libraries for low cost citizen science tools


Validate Citizen Science Tools





Support Citizen Science Research





Lending Libraries for Citizen Science Kits



ALL NEW Pole Mapping!



Balloon Mapping




Community-led Incubated Kits 2 items



DIY Spectrometry 9 items



Gifts & donations 3 items



Infrared Photography 5 items



Introductory Kits 9 items



Kite Mapping 13 items



Mapping Kits 5 items



Mapping Supplies
12 items



Publications 11 items



Wholesale 8 items

Creating an infrastructure for rigorous, responsive research that supports agencies and the public.

Open Source Research Tools-- low cost, DIY tools, that are in the public domain

Open Source Software--Publicly accessible databases and maps that enable rapid grassroots reporting and analysis across communities

- •Study Large Scale Industries and Shared Environmental Health Problems
- •From Community to Community
- Fuse Community learning/Advocacy and Academic/Regulatory Analysis
- And Improve Monitoring while Facilitating Basic Research on previously Inaccessible Problems

Emerging Health Threats

Health
Disparities/CBPR

Interdisciplinary Collaboration Scientific Literacy

Thank you!

- Research Team
- -Caitlin Kennedy (Drew University)
- -Megan McLaughlin (RISD)
- -Elisabeth Wilder (Northeastern University)
- -Hannah Gartner (Northeastern University)
- Community Partners and Collaborators
- -Clark Resource Council
- -Powder River Basin Resource Council
- -San Juan Citizen's Alliance
- SSEHRI/Northeastern University
- Public Lab (<u>www.publiclab).org</u>)
- ShaleTest.org





